

Errata for ICD-10-CM and ICD-10-PCS Coding Handbook 2019 with Answers

Narrative changes appear in bold italicized text; deletions show as strikethrough text.

8/29/18

Page Number	Current Text	Revision
329	Assign an additional code to identify the causal organism with codes in category O86.	Assign an additional code to identify the causal organism with codes in category O86. [New paragraph:] For sepsis following an obstetrical procedure, such as cesarean delivery, assign first a code from O86.00 to O86.03, Infection of obstetric surgical wound, that identifies the site of the infection (e.g. deep incisional site), if known. Assign an additional code for sepsis following an obstetrical procedure (O86.04). Use an additional code to identify the infectious agent (e.g. MRSA).
403	If the provider has specifically documented different causes for the hypertension and the heart condition, code the heart condition (I50.-, I51.4–I51.9) and hypertension separately.	If the provider has specifically documented different causes for the hypertension and the heart condition, code the heart condition (I50.-, I51.4– I51.9 I51.7, I51.89, I51.9) and hypertension separately.
408	However, in cases involving adverse effects of drugs, the sequencing of secondary pulmonary hypertension and associated conditions is based on the reason for the encounter.	However, in cases involving adverse effects of drugs, the sequencing of secondary pulmonary hypertension and associated conditions is based on the reason for the encounter. The sequencing is based on the reason for the encounter, except for adverse effects of drugs.
641	14. . . Postoperatively, the patient did well on antibiotics. . . . 10D00Z1 Extraction of products of conception, low cervical, open approach	14. . . Postoperatively, the patient did well on antibiotics. . . . 10D00Z1 Extraction of products of conception, low cervical , open approach